

\*c1cc(\*)c(\*)c2c1c(n2)C3=CC=CC=C3

1. (Original): The compound of the general formula (1):

W, X and Y are all CH and Z is N; R is halo;

$R^1$  is aryl, heteroaryl, morpholino, piperidino or pyrrolidino;  $R^2$  is  $NR^3R^4$ ,

R<sup>3</sup> and R<sup>4</sup> are independently H, C<sub>1-8</sub> alkyl, C<sub>2-8</sub> alkenyl, C<sub>2-8</sub> alkynyl, aryl, aryl(C<sub>1-8</sub>)alkyl, C<sub>3-8</sub> cycloalkyl, C<sub>3-8</sub> cycloalkyl(C<sub>1-6</sub>)alkyl, heteroaryl, heteroaryl(C<sub>1-8</sub>)alkyl, NR<sup>5</sup>R<sup>6</sup>, provided that not both R<sup>3</sup> and R<sup>4</sup> are H or NR<sup>5</sup>R<sup>6</sup>, or

R<sup>3</sup> and R<sup>4</sup> together form a C<sub>3-7</sub> alkylene or C<sub>3-7</sub> alkenylene chain optionally substituted with one or more C<sub>1-4</sub> alkyl or C<sub>1-4</sub> alkoxy groups, or,

together with the nitrogen atom to which they are attached, R<sup>3</sup> and R<sup>4</sup> form a morpholine, thiomorpholine, thiomorpholine S-oxide or thiomorpholine S-dioxide ring or a piperazine or piperazine *N*-(C<sub>1-4</sub>)alkyl (especially *N*-methyl) ring; and

R<sup>5</sup> and R<sup>6</sup> are independently H, C<sub>1-8</sub> alkyl, C<sub>2-8</sub> alkenyl, C<sub>2-8</sub> alkynyl, aryl, aryl(C<sub>1-8</sub>)alkyl, C<sub>3-8</sub> cycloalkyl, C<sub>3-8</sub> cycloalkyl(C<sub>1-6</sub>)alkyl, heteroaryl or hetero-aryl(C<sub>1-8</sub>)alkyl;

any of the foregoing alkyl, alkenyl, alkynyl or cycloalkyl groups or moieties (other than for R<sup>8</sup>) being optionally substituted with halogen, cyano, C<sub>1-6</sub> alkoxy, C<sub>1-6</sub> alkylcarbonyl, C<sub>1-6</sub> alkoxycarbonyl, C<sub>1-6</sub> haloalkoxy, C<sub>1-6</sub> alkylthio, tri(C<sub>1-4</sub>)alkylsilyl, C<sub>1-6</sub>alkylamino or C<sub>1-6</sub> dialkylamino.

any of the foregoing morpholine, thiomorpholine, piperidine, piperazine and pyrrolidine rings being optionally substituted with C<sub>1-4</sub> alkyl (especially methyl), and

any of the foregoing aryl or heteroaryl groups or moieties being optionally substituted with one or more substituents selected from halo, hydroxy, mercapto, C<sub>1-6</sub> alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>1-6</sub> alkoxy, C<sub>2-6</sub> alkenyloxy, C<sub>2-6</sub> alkynyloxy, halo(C<sub>1-6</sub>)alkyl, halo(C<sub>1-6</sub>)alkoxy, C<sub>1-6</sub> alkylthio, halo(C<sub>1-6</sub>)alkylthio, hydroxy(C<sub>1-6</sub>)alkyl, C<sub>1-4</sub>alkoxy(C<sub>1-6</sub>)alkyl, C<sub>3-6</sub> cycloalkyl, C<sub>3-6</sub> cycloalkyl(C<sub>1-4</sub>)alkyl, phenoxy, benzyloxy, benzoyloxy, cyano, isocyano, thiocyanato, isothiocyanato, nitro, -NR<sup>'''</sup>R<sup>'''</sup>, -NHCOR<sup>'''</sup>, -NHCONR<sup>'''</sup>R<sup>'''</sup>, -CONR<sup>'''</sup>R<sup>'''</sup>, -SO<sub>2</sub>R<sup>'''</sup>, -OSO<sub>2</sub>R<sup>'''</sup>, -COR<sup>'''</sup>, -CR<sup>'''</sup>=NR<sup>'''</sup> or -N=CR<sup>'''</sup>R<sup>'''</sup>, in which R<sup>'''</sup> and R<sup>'''</sup> are independently hydrogen, C<sub>1-4</sub> alkyl, halo(C<sub>1-4</sub>)alkyl, C<sub>1-4</sub> alkoxy, halo(C<sub>1-4</sub>)alkoxy, C<sub>1-4</sub> alkylthio, C<sub>3-6</sub> cycloalkyl, C<sub>3-6</sub> cycloalkyl(C<sub>1-4</sub>)alkyl, phenyl or benzyl, the phenyl and benzyl groups being optionally substituted with halogen, C<sub>1-4</sub> alkyl or C<sub>1-4</sub> alkoxy.

2. (Original): A compound according to claim 1 wherein  
 $R^3$  is  $C_{1-8}$  alkyl, halo( $C_{1-8}$ )alkyl, hydroxy( $C_{1-8}$ )alkyl,  $C_{1-4}$  alkoxy( $C_{1-8}$ )alkyl,  $C_{1-4}$  alkoxyhalo( $C_{1-8}$ )alkyl, tri( $C_{1-4}$ )alkylsilyl( $C_{1-6}$ )alkyl,  $C_{1-4}$  alkylcarbonyl( $C_{1-8}$ )alkyl,  $C_{1-4}$  alkylcarbonylhalo( $C_{1-8}$ )alkyl, phenyl( $C_{1-4}$ )alkyl,  $C_{2-8}$  alkenyl, halo( $C_{2-8}$ )alkenyl,  $C_{2-8}$  alkynyl,  $C_{3-8}$  cycloalkyl optionally substituted with chloro, fluoro or methyl,  $C_{3-8}$  cycloalkyl-( $C_{1-4}$ )alkyl, phenylamino, piperidino or morpholino, the phenyl ring of phenylalkyl or phenylamino being optionally substituted with one, two or three substituents selected from halo,  $C_{1-4}$  alkyl, halo( $C_{1-4}$ )alkyl,  $C_{1-4}$  alkoxy and halo( $C_{1-4}$ )alkoxy; and  
 $R^4$  is H,  $C_{1-4}$  alkyl, halo( $C_{1-4}$ )alkyl or amino, or  
 $R^3$  and  $R^4$  together form a  $C_{3-7}$  alkylene or  $C_{3-7}$  alkenylene chain optionally substituted with methyl, or,  
together with the nitrogen atom to which they are attached,  $R^3$  and  $R^4$  form a morpholine, thiomorpholine, thiomorpholine S-oxide or thiomorpholine S-dioxide ring or a piperazine or piperazine *N*-( $C_{1-4}$ )alkyl (especially *N*-methyl) ring, in which the morpholine or piperazine rings are optionally substituted with methyl.
3. (Currently Amended): A compound according to claim 1 ~~or 2~~ claims wherein  
 $R^1$  is phenyl optionally substituted with from one to five halogen atoms or with from one to three substituents selected from halo,  $C_{1-4}$  alkyl, halo( $C_{1-4}$ )alkyl,  $C_{1-4}$  alkoxy or halo( $C_{1-4}$ )-alkoxy, pyridyl optionally substituted with from one to four halogen atoms or with from one to three substituents selected from halo,  $C_{1-4}$  alkyl, halo( $C_{1-4}$ )alkyl,  $C_{1-4}$  alkoxy or halo( $C_{1-4}$ )-alkoxy, 2- or 3-thienyl optionally substituted with from one to three halogen atoms or with from one to three substituents selected from halo,  $C_{1-4}$ alkyl, halo( $C_{1-4}$ )alkyl,  $C_{1-4}$  alkoxy or halo( $C_{1-4}$ )alkoxy, or piperidino or morpholino both optionally substituted with one or two methyl groups.
4. (Original): A compound according to claim 3 wherein  $R^1$  is 2,6-difluorophenyl, 2-fluoro-6-chlorophenyl, 2,5,6-trifluorophenyl, 2,4,6-trifluorophenyl, 2,6-difluoro-4-methoxyphenyl or pentafluorophenyl.
5. (Original): A compound according to claim 1 wherein W, X and Y are all CH and Z is N; R is halo,  $R^1$  is aryl, heteroaryl, morpholino, piperidino or pyrrolidino;  $R^2$  is  $NR^3R^4$ ;  $R^3$  and  $R^4$  are independently H,  $C_{1-8}$  alkyl,  $C_{2-8}$  alkenyl,  $C_{2-8}$  alkynyl, aryl, aryl( $C_{1-8}$ )alkyl,  $C_{3-8}$  cycloalkyl,  $C_{3-8}$  cycloalkyl( $C_{1-6}$ )alkyl, heteroaryl, heteroaryl( $C_{1-8}$ )alkyl,  $NR^5R^6$ , provided that

not both  $R^3$  and  $R^4$  are H or  $NR^5R^6$ , or

$R^3$  and  $R^4$  together form a  $C_{3-7}$  alkylene or  $C_{3-7}$  alkenylene chain optionally substituted with one or more  $C_{1-4}$  alkyl or  $C_{1-4}$  alkoxy groups, or,

together with the nitrogen atom to which they are attached,  $R^3$  and  $R^4$  form a morpholine, thiomorpholine, thiomorpholine S-oxide or thiomorpholine S-dioxide ring or a piperazine or piperazine  $N$ -( $C_{1-4}$ )alkyl (especially  $N$ -methyl) ring; and

$R^5$  and  $R^6$  are independently H,  $C_{1-8}$  alkyl,  $C_{2-8}$  alkenyl,  $C_{2-8}$  alkynyl, aryl, aryl( $C_{1-8}$ )alkyl,  $C_{3-8}$  cycloalkyl,  $C_{3-8}$  cycloalkyl( $C_{1-6}$ )alkyl, heteroaryl or heteroaryl( $C_{1-8}$ )alkyl;

any of the foregoing alkyl, alkenyl, alkynyl or cycloalkyl groups or moieties being optionally substituted with halogen, cyano,  $C_{1-6}$ alkoxy,  $C_{1-6}$  alkylcarbonyl,  $C_{1-6}$  alkoxy carbonyl,  $C_{1-6}$  haloalkoxy,  $C_{1-6}$  alkylthio, tri( $C_{1-4}$ )alkylsilyl,  $C_{1-6}$  alkylamino or  $C_{1-6}$  dialkylamino, any of the foregoing morpholine, thiomorpholine, piperidine, piperazine and pyrrolidine rings being optionally substituted with  $C_{1-4}$  alkyl (especially methyl), and any of the foregoing aryl, heteroaryl, aryloxy or heteroaryl groups being optionally substituted with one or more substituents selected from halo, hydroxy, mercapto,  $C_{1-6}$  alkyl,  $C_{2-6}$  alkenyl,  $C_{2-6}$  alkynyl,  $C_{1-6}$  alkoxy,  $C_{2-6}$  alkenyloxy,  $C_{2-6}$  alkynyloxy, halo( $C_{1-6}$ )alkyl, halo( $C_{1-6}$ )alkoxy,  $C_{1-6}$  alkylthio, halo( $C_{1-6}$ )alkylthio, hydroxy( $C_{1-6}$ )alkyl,  $C_{1-4}$  alkoxy( $C_{1-6}$ )alkyl,  $C_{3-6}$  cycloalkyl,  $C_{3-6}$  cycloalkyl( $C_{1-4}$ )alkyl, phenoxy, benzyloxy, benzoyloxy, cyano, isocyano, thiocyanato, isothiocyanato, nitro,  $-NR''R'''$ ,  $-NHCOR''$ ,  $-NHCONR''R'''$ ,  $-CONR''R'''$ ,  $-SO_2R''$ ,  $-OSO_2R''$ ,  $-COR''$ ,  $-CR''=NR'''$  or  $-N=CR''R'''$ , in which  $R''$  and  $R'''$  are independently hydrogen,  $C_{1-4}$ alkyl, halo( $C_{1-4}$ )alkyl,  $C_{1-4}$  alkoxy, halo( $C_{1-4}$ )alkoxy,  $C_{1-4}$  alkylthio,  $C_{3-6}$  cycloalkyl,  $C_{3-6}$  cycloalkyl( $C_{1-4}$ )alkyl, phenyl or benzyl, the phenyl and benzyl groups being optionally substituted with halogen,  $C_{1-4}$  alkyl or  $C_{1-4}$  alkoxy.

6. (Original): A compound according to claim 1 wherein W, X and Y are all CH and Z is N; R is halo;  $R^1$  is aryl, heteroaryl, morpholino, piperidino or pyrrolidino;  $R^2$  is  $NR^3R^4$ ,  $R^3$  is  $C_{1-4}$  alkyl, halo( $C_{1-4}$ )alkyl,  $C_{2-4}$  alkenyl,  $C_{3-6}$  cycloalkyl,  $C_{3-6}$  cycloalkyl( $C_{1-4}$ )alkyl or phenylamino in which the phenyl ring is optionally substituted with one, two or three substituents selected from halo,  $C_{1-4}$  alkyl, halo( $C_{1-4}$ )alkyl,  $C_{1-4}$  alkoxy and halo( $C_{1-4}$ )alkoxy; and  $R^4$  is H,  $C_{1-4}$  alkyl or amino, or  $R^3$  and  $R^4$  together form a  $C_{4-6}$  alkylene chain optionally substituted with  $C_{1-4}$  alkyl or  $C_{1-4}$  alkoxy, or together with the nitrogen atom to which they are attached,  $R^3$  and  $R^4$  form a morpholine, thiomorpholine, thiomorpholine S-oxide or thiomorpholine S-dioxide ring or a piperazine or piperazine  $N$ -( $C_{1-4}$ )alkyl (especially  $N$ -methyl) ring;

any of the foregoing alkyl, alkenyl, alkynyl or cycloalkyl groups or moieties being optionally substituted with halogen, cyano, C<sub>1-6</sub> alkoxy, C<sub>1-6</sub>alkylcarbonyl, C<sub>1-6</sub>alkoxycarbonyl, C<sub>1-6</sub> haloalkoxy, C<sub>1-6</sub> alkylthio, tri(C<sub>1-4</sub>)alkylsilyl, C<sub>1-6</sub> alkylamino or C<sub>1-6</sub>dialkylamino, any of the foregoing morpholine, thiomorpholine, piperidine, piperazine and pyrrolidine rings being optionally substituted with C<sub>1-4</sub> alkyl (especially methyl), and any of the foregoing aryl or heteroaryl groups or moieties being optionally substituted with one or more substituents selected from halo, hydroxy, mercapto, C<sub>1-6</sub> alkyl, C<sub>2-6</sub> alkenyl, C<sub>2-6</sub> alkynyl, C<sub>1-6</sub> alkoxy, C<sub>2-6</sub> alkenyloxy, C<sub>2-6</sub>alkynyloxy, halo(C<sub>1-6</sub>)alkyl, halo(C<sub>1-6</sub>)alkoxy, C<sub>1-6</sub> alkylthio, halo(C<sub>1-6</sub>)alkylthio, hydroxy(C<sub>1-6</sub>)alkyl, C<sub>1-4</sub> alkoxy(C<sub>1-6</sub>)alkyl, C<sub>3-6</sub> cycloalkyl, C<sub>3-6</sub> cycloalkyl(C<sub>1-4</sub>)alkyl, phenoxy, benzyloxy, benzoyloxy, cyano, isocyano, thiocyanato, isothiocyanato, nitro, -NR<sup>'''</sup>R<sup>'''</sup>, -NHCOR<sup>'''</sup>, -NHCONR<sup>'''</sup>R<sup>'''</sup>, -CONR<sup>'''</sup>R<sup>'''</sup>, -SO<sub>2</sub>R<sup>'''</sup>, -OSO<sub>2</sub>R<sup>'''</sup>, -COR<sup>'''</sup>, -CR<sup>'''</sup>=NR<sup>'''</sup> or -N=CR<sup>'''</sup>R<sup>'''</sup>, in which R<sup>'''</sup> and R<sup>'''</sup> are independently hydrogen, C<sub>1-4</sub>alkyl, halo(C<sub>1-4</sub>)alkyl, C<sub>1-4</sub> alkoxy, halo(C<sub>1-4</sub>)alkoxy, C<sub>1-4</sub> alkylthio, C<sub>3-6</sub> cycloalkyl, C<sub>3-6</sub> cycloalkyl(C<sub>1-4</sub>)alkyl, phenyl or benzyl, the phenyl and benzyl groups being optionally substituted with halogen, C<sub>1-4</sub> alkyl or C<sub>1-4</sub> alkoxy.

7. (Original): A compound according to claim 1 wherein one of W, X and Y are all CH and Z is N;

R is halo; R<sup>1</sup> is optionally substituted phenyl; R<sup>2</sup> is NR<sup>3</sup>R<sup>4</sup>,

R<sup>3</sup> and R<sup>4</sup> are independently H, C<sub>1-8</sub> alkyl, C<sub>2-8</sub> alkenyl, C<sub>2-8</sub> alkynyl, aryl, aryl(C<sub>1-8</sub>)alkyl, C<sub>3-8</sub> cycloalkyl, C<sub>3-8</sub> cycloalkyl(C<sub>1-6</sub>)alkyl, heteroaryl, heteroaryl(C<sub>1-8</sub>)alkyl, NR<sup>5</sup>R<sup>6</sup>, provided that not both R<sup>3</sup> and R<sup>4</sup> are H or NR<sup>5</sup>R<sup>6</sup>, or

R<sup>3</sup> and R<sup>4</sup> together form a C<sub>3-7</sub> alkylene or C<sub>3-7</sub> alkenylene chain optionally substituted with one or more C<sub>1-4</sub> alkyl or C<sub>1-4</sub> alkoxy groups, or,

together with the nitrogen atom to which they are attached, R<sup>3</sup> and R<sup>4</sup> form a morpholine, thiomorpholine, thiomorpholine S-oxide or thiomorpholine S-dioxide ring or a piperazine or piperazine N-(C<sub>1-4</sub>)alkyl (especially N-methyl) ring; and



R<sup>5</sup> and R<sup>6</sup> are independently H, C<sub>1-8</sub> alkyl, C<sub>2-8</sub> alkenyl, C<sub>2-8</sub> alkynyl, aryl, aryl(C<sub>1-8</sub>)alkyl, C<sub>3-8</sub> cycloalkyl, C<sub>3-8</sub> cycloalkyl(C<sub>1-6</sub>)alkyl, heteroaryl or heteroaryl(C<sub>1-8</sub>)alkyl;

any of the foregoing alkyl, alkenyl, alkynyl or cycloalkyl groups or moieties being optionally substituted with halogen, cyano, C<sub>1-6</sub> alkoxy, C<sub>1-6</sub> alkylcarbonyl, C<sub>1-6</sub>alkoxycarbonyl, C<sub>1-6</sub> haloalkoxy, C<sub>1-6</sub> alkylthio, tri(C<sub>1-4</sub>)alkylsilyl, C<sub>1-6</sub> alkylamino or C<sub>1-6</sub>dialkylamino; any of the foregoing morpholine, thiomorpholine, piperidine, piperazine and pyrrolidine rings being optionally substituted with C<sub>1-4</sub> alkyl (especially methyl), and any of the foregoing aryl or heteroaryl groups or moieties, including the phenyl group of R<sup>1</sup>, being optionally substituted

with one or more substituents selected from halo, hydroxy, mercapto, C<sub>1-6</sub> alkyl, C<sub>2-6</sub> alkenyl, C<sub>2-6</sub> alkynyl, C<sub>1-6</sub>alkoxy, C<sub>2-6</sub> alkenyloxy, C<sub>2-6</sub> alkynyloxy, halo(C<sub>1-6</sub>)alkyl, halo(C<sub>1-6</sub>)alkoxy, C<sub>1-6</sub>alkylthio, halo(C<sub>1-6</sub>)alkylthio, hydroxy(C<sub>1-6</sub>)alkyl, C<sub>1-4</sub> alkoxy(C<sub>1-6</sub>)alkyl, C<sub>3-6</sub>cycloalkyl, C<sub>3-6</sub> cycloalkyl(C<sub>1-4</sub>)alkyl, phenoxy, benzyloxy, benzoyloxy, cyano, isocyano, thiocyanato, isothiocyanato, nitro, -NR<sup>'''</sup>R<sup>'''</sup>, -NHCOR<sup>'''</sup>, -NHCONR<sup>'''</sup>R<sup>'''</sup>, -CONR<sup>'''</sup>R<sup>'''</sup>, -SO<sub>2</sub>R<sup>'''</sup>, -OSO<sub>2</sub>R<sup>'''</sup>, -COR<sup>'''</sup>, -CR<sup>'''</sup>=NR<sup>'''</sup> or -N=CR<sup>'''</sup>R<sup>'''</sup>, in which R<sup>'''</sup> and R<sup>'''</sup> are independently hydrogen, C<sub>1-4</sub> alkyl, halo(C<sub>1-4</sub>)alkyl, C<sub>1-4</sub> alkoxy, halo(C<sub>1-4</sub>)alkoxy, C<sub>1-4</sub> alkylthio, C<sub>3-6</sub> cycloalkyl, C<sub>3-6</sub> cycloalkyl(C<sub>1-4</sub>)alkyl, phenyl or benzyl, the phenyl and benzyl groups being optionally substituted with halogen, C<sub>1-4</sub>alkyl or C<sub>1-4</sub> alkoxy.

8. (Original): A compound according to claim 1 wherein W, X and Y are all CH and Z is N; R is halo; R<sup>1</sup> is phenyl optionally substituted with from one to five halogen atoms or with from one to three substituents selected from halo, C<sub>1-4</sub> alkyl, halo(C<sub>1-4</sub>)alkyl, C<sub>1-4</sub>alkoxy or halo(C<sub>1-4</sub>)-alkoxy, pyridyl optionally substituted with from one to four halogen atoms or with from one to three substituents selected from halo, C<sub>1-4</sub> alkyl, halo(C<sub>1-4</sub>)alkyl, C<sub>1-4</sub>alkoxy or halo(C<sub>1-4</sub>)-alkoxy, 2- or 3-thienyl optionally substituted with from one to three halogen atoms or with from one to three substituents selected from halo, C<sub>1-4</sub>alkyl, halo(C<sub>1-4</sub>)alkyl, C<sub>1-4</sub> alkoxy or halo(C<sub>1-4</sub>)alkoxy, or piperidino or morpholino both optionally substituted with one or two methyl groups; R<sup>2</sup> is NR<sup>3</sup>R<sup>4</sup>;
- R<sup>3</sup> is C<sub>1-8</sub> alkyl, halo(C<sub>1-8</sub>)alkyl, hydroxy(C<sub>1-8</sub>)alkyl, C<sub>1-4</sub> alkoxy(C<sub>1-8</sub>)alkyl, C<sub>1-4</sub> alkoxyhalo(C<sub>1-8</sub>)alkyl, tri(C<sub>1-4</sub>)alkylsilyl(C<sub>1-6</sub>)alkyl, C<sub>1-4</sub> alkylcarbonyl(C<sub>1-8</sub>)alkyl, C<sub>1-4</sub> alkylcarbonylhalo(C<sub>1-8</sub>)alkyl, phenyl(C<sub>1-4</sub>)alkyl, C<sub>2-8</sub> alkenyl, halo(C<sub>2-8</sub>)alkenyl, C<sub>2-8</sub> alkynyl, C<sub>3-8</sub> cycloalkyl optionally substituted with chloro, fluoro or methyl, C<sub>3-8</sub> cycloalkyl(C<sub>1-4</sub>)alkyl, phenylamino, piperidino or morpholino, the phenyl ring of phenylalkyl or phenylamino being optionally substituted with one, two or three substituents selected from halo, C<sub>1-4</sub> alkyl, halo(C<sub>1-4</sub>)alkyl, C<sub>1-4</sub> alkoxy and halo(C<sub>1-4</sub>)alkoxy; and
- R<sup>4</sup> is H, C<sub>1-4</sub> alkyl, halo(C<sub>1-4</sub>)alkyl or amino, or
- R<sup>3</sup> and R<sup>4</sup> together form a C<sub>3-7</sub> alkylene or C<sub>3-7</sub> alkenylene chain optionally substituted with methyl, or,
- together with the nitrogen atom to which they are attached, R<sup>3</sup> and R<sup>4</sup> form a morpholine, thiomorpholine, thiomorpholine S-oxide or thiomorpholine S-dioxide ring or a piperazine or piperazine N-(C<sub>1-4</sub>)alkyl (especially N-methyl) ring, in which the morpholine or piperazine rings are optionally substituted with methyl.

9. (Original): A compound according to claim 1 wherein one of W, X and Y are all CH and Z is N;  
 R is halo; R<sup>1</sup> is phenyl optionally substituted with from one to five halogen atoms or with from one to three substituents selected from halo, C<sub>1-4</sub> alkyl, halo(C<sub>1-4</sub>)alkyl, C<sub>1-4</sub>alkoxy or halo(C<sub>1-4</sub>)alkoxy; R<sup>2</sup> is NR<sup>3</sup>R<sup>4</sup>; R<sup>3</sup> is C<sub>1-4</sub> alkyl, halo(C<sub>1-4</sub>)alkyl, C<sub>2-4</sub>alkenyl, C<sub>3-6</sub> cycloalkyl, C<sub>3-6</sub> cycloalkyl(C<sub>1-4</sub>)alkyl or phenylamino in which the phenyl ring is optionally substituted with one, two or three substituents selected from halo, C<sub>1-4</sub> alkyl, halo(C<sub>1-4</sub>)alkyl, C<sub>1-4</sub> alkoxy and halo(C<sub>1-4</sub>)alkoxy; and  
 R<sup>4</sup> is H, C<sub>1-4</sub> alkyl or amino, or R<sup>3</sup> and R<sup>4</sup> together form a C<sub>4-6</sub> alkylene chain optionally substituted with methyl, or, together with the nitrogen atom to which they are attached, R<sup>3</sup> and R<sup>4</sup> form a morpholine ring.
10. (Original): A process for preparing a compound of the general formula (1) according to claim 1 wherein one of R is chloro or fluoro and R<sup>2</sup> is NR<sup>3</sup>R<sup>4</sup> and W, X, Y, Z, R<sup>1</sup>, R<sup>3</sup> and R<sup>4</sup> are as defined in claim 1, which comprises reacting an amine of the general formula NR<sup>3</sup>R<sup>4</sup> with a compound of the general formula (6) or (13):



wherein W, X, Y, Z and R<sup>1</sup> are as defined in claim 1.

11. (Original): A plant fungicidal composition comprising a fungicidally effective amount of a compound as defined in claim 1 and a suitable carrier or diluent therefor.
12. (Currently Amendment): A method of combating or controlling phytopathogenic fungi which comprises applying to a plant, to a seed of a plant, to the locus of the plant or seed or to soil or to any other plant growth medium, a fungicidally effective amount of a compound according to claim 1 ~~or a composition according to claim 12.~~